

CERHR Update and Evaluation Concept:

State-of-the-Science Evaluation on Environmental Exposures and Diabetes/Obesity

Kristina Thayer, Ph.D.

National Institute of Environmental Health Sciences

NTP Board of Scientific Counselors Meeting December 9, 2009





Outline

- New CERHR staff
- Updated evaluation of soy formula
- Concept proposal for state of the science assessment of environmental contaminants and diabetes/obesity



New CERHR Staff

- 4 new staff since August 2009
 - Vickie Walker, BS
 - Kyla Taylor, MS
 - Andy Rooney, Ph.D
 - Kembra Howdeshell, Ph.D
- Total staff of 6
 - 4 new staff, Mike Shelby, Kris Thayer



Soy Formula Evaluation

- October 19, 2009: Release draft Expert Panel Report
- December 16 18, 2009: Expert Panel meeting
- Early January 2010: Release final Expert Panel report for public comment
- March 2010: Release Draft NTP Brief on Soy Formula for public comment
- May 10, 2010: Peer review of draft NTP Brief by BSC
- Summer 2010: Release final NTP Monograph on Soy Formula





Concept Proposal

State-of-the-science evaluation of environmental exposures and diabetes/obesity





Justification

- Diabetes and obesity are major threats to human health
 - ~40% of people ≥20 of age have diabetes or pre-diabetes, NHANES 2005-2006 (Cowie 2009)
 - 12.9% diabetes (of which ~40% undiagnosed); 29.5% with pre-diabetes (impaired fasting glucose or impaired 2-hour glucose tolerance)
 - − ~70% of type 2 diabetes risk attributed to overweight/obesity (Eyre 2004),
 - Thus, ~30% not accounted for by body weight
 - Often discussed in the context of developmental origins of adult disease or prenatal programming, e.g., PPTOXII, Miami December 7-10, 2009
 - Risk factors for impaired reproduction/development



Justification

- Growing scientific and public interest
 - Organohalogens: ~ 40 human studies since 2005 (~70 total)
 - Arsenic: Odds ratio of 3.58 (95%CI: 1.18-10.83) for type 2 diabetes based on comparison of 80th and 20th percentiles of total urinary arsenic, 2003-2004 NHANES data (Navas-Acien 2008)
 - BPA: Higher urinary levels associated with diabetes and cardiovascular disease, 2003-2004 NHANES data (Lang 2008)
 - Diabetes: Odds ratio per 1-SD increase in BPA concentration = 1.39 (95% CI: 1.21-1.60)
 - Newsweek, September 11, 2009 "Born to be Big: Early Exposure to Common Chemicals may be Programming Kids to be Fat"



Goals and Expected Outcomes

Charge 1: Critical assessment of the current literature

- a) Evaluate strength/weaknesses, consistency, and biological plausibility of findings reported in humans and experimental animals within and across chemicals;
 - NOT intended to develop "level of concern" conclusions
- b) Identify the most useful and relevant endpoints in experimental animals and *in vitro* models

Charge 2: Focus future research directions

a) Identify data gaps and areas for future evaluation/research



Proposed Approach - Workshop

Present concept to NTP Board of Scientific Counselors

CERHR staff prepare literature review document

Identify experts and convene subpanels of scientists to address charge #1 (critical assessment of current literature) for each exposure chapter

Public Comment

Public Workshop (early Fall 2010)

Oral public comments

Subpanels finalize responses to charge #1, present in plenary

· Group discussion on subpanel findings`

Second round of subpanels meet to address charge #2 (data gaps and research needs)

• Group discussion on research needs

Open general discussion

Prepare NTP Monograph (late Fall 2010)

Workshop report as executive summary + literature review document (workshop report also submitted for publication)



Literature Screen

- Use of a combination of health- and exposure-based MeSH terms
- Iterative process
 - Wish list: broad screen of environmental exposures for effects related to diabetes, metabolic syndrome, and obesity
 - A front end screen for health-effects related to metabolic syndrome not feasible, i.e., too many articles retrieved
 - Can be follow-up search for specific exposures
 - "Obesity" not a precise MeSH heading for this evaluation
 - · Targeted approach
 - MeSH terms for environmental exposures too broad*
 - · Focus on specific exposures or classes of chemicals

^{*}MeSH headings: environmental pollution, toxic action, endocrine disruptors



Extent of Literature

- Nature of literature varies from exposure to exposure, i.e., number of publications, proportion of human/animal/in vitro
 - Relatively large literatures
 - Arsenic, organochlorines (e.g., dioxin, PCBs, DDT/DDE), genistein
 - Relatively small literature
 - Organotins, BPA, phthalates
- Thus, the type of conclusions reached for each type of exposure will vary
 - Range from research focusing to those that permit drawing conclusions on consistency/biological plausibility




Approach to Assess "Obesogens"

- Targeted search
 - e.g., tributyltin, genistein, BPA, DEHP, PFOA
 - Consider consistency of effects on body weight following developmental exposure
 - BUT, recognize that topic is more complex than effects on body weight




Tributyltin as Inducer of Adipogenesis (Grun et al., 2006)


- C57BL/6 dams treated with 0.5 mg/kg TBT by ip injection on GD12-18; looked at effects in F1 offspring:
 - ↑ lipid accumulation in adipose depots, liver and testis in neonates
 - † epididymal adipose mass in adult male offspring
 - No effects on body weight in adult mice



Control TBT

Mammary adipose tissue in neonates
(Oil red O stain)







Type of Scientific Expertise

- Charge 1: Critical assessment of the current literature
 - Include expertise in particular exposures as well as expertise in the primary health conditions
- Charge 2: Focus future research directions
 - Overlap with 1st subpanel membership, but supplemented with others to generate both general research suggestions as well as those specific to NIEHS/NTP



Charge to Board

To review and comment on proposed CERHR evaluation concept and determine whether the evaluation is an appropriate use of NTP program resources.

- Rationale
- Merit
- Overall significance and public health impact
- Scope of review activity
- Other comments



References

- Cowie *et al.*, 2009. Full accounting of diabetes and pre-diabetes in the US population in 1988-1994 and 2005-2006. Diabetes Care. 32: 287-294
- Dunnick *et al.*, 2007. Inclusion of biomarkers for detecting perturbations in the heart and lung and lipid/carbohydrate metabolism in National Toxicology Program studies. Toxicol Sci. 2007 Nov;100(1):29-35.
- Eyre et al., 2004. Preventing cancer, cardiovascular disease, and diabetes: A common agenda for the American Cancer Society, the American Diabetes Association, and the American Heart Association. CA Cancer J Clin 54: 190-2007
- Grun et al., 2006. Endocrine-disrupting organotin compounds are potent inducers of adipogenesis in vertebrates. Mol Endocrinol. 20(9):2141-55.
- Lang et al., 2008. Association of urinary bisphenol A concentration with medical disorders and laboratory abnormalities in adults. Jama. 300:1303-1310.
- Navas-Acien *et al.*, 2008. Arsenic exposure and prevalence of type 2 diabetes in US adults. Jama. 300:814-822.
- Ohlsson et al., 2000. Obesity and disturbed lipoprotein profile in estrogen receptor-alphadeficient male mice. Biochem Biophys Res Commun. 278(3):640-5.